

# **Anti-Oct-2 Rabbit Monoclonal Antibody**

## Introduction

Transcription factor that specifically binds to the octamer motif (5'-ATTTGCAT-3'). Regulates transcription in a number of tissues in addition to activating immunoglobulin gene expression.

## Product parameters

Alternative Names	POU2F2; OCT2; OTF2; POU domain; class 2; transcription factor 2; Lymphoid-restricted immunoglobuli octamer-binding protein NF-A2; Octamer-binding protein 2; Oct-2; Octamer-binding transcription factor 2
, mornauvo riameo	OTF-2
Gene ID	5452
Gene Name	POU2F2
SwissProt ID	P09086
Host	Rabbit
Reactivity	Rat
Molecular Weight	Calculated MW: 51 kDa; Observed MW: 55 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-17F9D4
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	A synthetic peptide of human Oct-2
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB
Dilution Ratio	WB: 1/500-1/1000
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Shipping	Blue ice

Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

#### Protocol

Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

#### Note

1. This product is for scientific research use only.





