

Anti-Phospho-DNA PKcs (Ser2056) Rabbit Monoclonal Antibody

Introduction

The PRKDC gene encodes the catalytic subunit of a nuclear DNA-dependent serine/threonine protein kinase (DNA-PK). The second component is the autoimmune antigen Ku (MIM 152690), which is encoded by the G22P1 gene on chromosome 22q. On its own, the catalytic subunit of DNA-PK is inactive and relies on the G22P1 component to direct it to the DNA and trigger its kinase activity; PRKDC must be bound to DNA to express its catalytic properties.

Product parameters

Alternative Names	PRKDC; HYRC1; DNA-dependent protein kinase catalytic subunit; DNA-PK catalytic subun DNA-PKcs; DNPK1; p460
Gene ID	5591
Gene Name	PRKDC
SwissProt ID	P78527
Host Achieve Perfect	Rabbit
Reactivity	Human
Molecular Weight	Calculated MW: 469 kDa; Observed MW: 469 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Phosphorylated
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-1H11B5
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human Phospho-DNA PKcs (S2056)
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P, ICC/IF
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200
Research Field	Epigenetics and Nuclear Signaling

Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.





Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





