

Anti-Phospho-Glycogen synthase (Ser641) Rabbit Monoclonal Antibody

Introduction

Transfers the glycosyl residue from UDP-Glc to the non-reducing end of alpha-1,4-glucan. Allosteric activation by glucose-6-phosphate. Phosphorylation reduces the activity towards UDP-glucose. When in the non-phosphorylated state, glycogen synthase does not require glucose-6-phosphate as an allosteric activator; when phosphorylated it does.

Product parameters

Alternative Names	GYS1; GYS; Glycogen [starch] synthase; muscle
Gene ID	2997
Gene Name	GYS1
SwissProt ID	P13807
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 84 kDa; Observed MW: 84 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Phosphorylated
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-5F1G3
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Ser641 of human Glycogen synthase 1/GYS1
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB, IHC-P, IP
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20
Research Field	Signal Transduction
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol



Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

