

## Anti-Phospho-Histone H1.4 (Thr17) Rabbit Monoclonal Antibody

### Introduction

Histone H1 protein binds to linker DNA between nucleosomes forming the macromolecular structure known as the chromatin fiber. Histones H1 are necessary for the condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation (By similarity).

### Product parameters

Alternative Names	Histone H1b; Histone H1s-4
Gene ID	3008
Gene Name	H1-4
SwissProt ID	P10412
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 22 kDa; Observed MW: 30 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Phosphorylated
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-16C2G9
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human Phospho-Histone H1.4 (T17)
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P, ICC/IF
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

## Protocol



Configure the product according to the application range and recommended dilution ratio.

**\*Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.



**APExBIO Technology**

**[www.apexbt.com](http://www.apexbt.com)**

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: [info@apexbt.com](mailto:info@apexbt.com)

