

Anti-Phospho-LAT (Tyr191) Rabbit Monoclonal Antibody

Introduction

Required for TCR (T-cell antigen receptor)- and pre-TCR-mediated signaling, both in mature T-cells and during their development. Involved in FCGR3 (low affinity immunoglobulin gamma Fc region receptor III)-mediated signaling in natural killer cells and FCER1 (high affinity immunoglobulin epsilon receptor)-mediated signaling in mast cells. Couples activation of these receptors and their associated kinases with distal intracellular events such as mobilization of intracellular calcium stores, PKC activation, MAPK activation or cytoskeletal reorganization through the recruitment of PLCG1, GRB2, GRAP2, and other signaling molecules.

Product parameters

| | |
|-------------------|--|
| Alternative Names | LAT; Linker for activation of T-cells family member 1; 36 kDa phospho-tyrosine adapter protein; pp36; p36-38 |
| Gene ID | 27040 |
| Gene Name | LAT |
| SwissProt ID | O43561 |
| Host | Rabbit |
| Reactivity | Human, Mouse, Rat |
| Molecular Weight | Calculated MW: 28 kDa; Observed MW: 38 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Phosphorylated |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-3G10A9 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Free |
| Immunogen | A synthesized peptide derived from human Phospho-LAT (Y191) |
| Purification | Affinity Chromatography |
| Buffer System | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Application | WB, IHC-P, ICC/IF, IP |
| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/30 |

| | |
|--------------------|----------------------------------|
| Research Field | Immunology |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |
| Expiration Date | 12 months |
| Note | Please avoid freeze-thaw cycles. |



Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

