

Anti-Phospho-Rac1/CDC42 (Ser71) Rabbit Monoclonal Antibody

Introduction

Rac and Cdc42 are members of the Rho-GTPase family. In mammals, Rac exists as three isoforms, Rac1, Rac2 and Rac3, which are highly similar in sequence. Rac1 and Cdc42, the most widely studied of this group, are ubiquitously expressed. Rac2 is expressed in cells of hematopoietic origin, and Rac3, while highly expressed in brain, is also found in many other tissues. Rac and Cdc42 play key signaling roles in cytoskeletal reorganization, membrane trafficking, transcriptional regulation, cell growth and development.

Product parameters

Alternative Names	CDC42; CDC42Hs; G25K; TKS; MIG5; Ras like protein TC25
Gene ID	998/5879
Gene Name	CDC42/RAC1
SwissProt ID	P60953/P63000
Host	Rabbit O
Reactivity	Human Achieve Perfection, Explore the Unknown
Molecular Weight	Calculated MW: 21 kDa; Observed MW: 24 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Phosphorylated
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-20G7H9
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Im <mark>muno</mark> gen	A synthesized peptide derived from human Phospho-Rac1/Cdc42 (Ser71)
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB
Dilution Ratio	WB: 1/500-1/1000
Research Field	Cell Biology

Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.





Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





