

Anti-Phospho-Tyrosine Rabbit Monoclonal Antibody

Introduction

Tyrosine phosphorylation plays a key role in cellular signaling. Research studies have shown that in cancer, unregulated tyrosine kinase activity can drive malignancy and tumor formation by generating inappropriate proliferation and survival signals. Antibodies specific for phospho-tyrosine have been invaluable reagents in these studies.

Product parameters

Alternative Names	-
Gene ID	-
Gene Name	-
SwissProt ID	-
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Refer to figures
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Phosphorylated
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-7B5B4
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from Phosphotyrosine
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, ICC/IF, FC, IP
Dilution Ratio	WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/50 FC: 1/50-1/100
Research Field	Neuroscience
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol



Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

