

Anti-PI3 Kinase p85 beta Rabbit Monoclonal Antibody

Introduction

Phosphoinositide 3-kinase (PI3K) catalyzes the production of phosphatidylinositol-3,4,5-triphosphate by phosphorylating phosphatidylinositol (PI), phosphatidylinositol-4-phosphate (PIP) and phosphatidylinositol-4,5-bisphosphate (PIP₂). Growth factors and hormones trigger this phosphorylation event, which in turn coordinates cell growth, cell cycle entry, cell migration, and cell survival.

Product parameters

Alternative Names	p85; p85 beta; p85-BETA; P85B; P85B_HUMAN; Phosphatidylinositol 3 kinase; Phosphatidylinositol 3 kinase regulatory beta subunit; Phosphatidylinositol 3 kinase regulatory subunit beta; Phosphatidylinositol 3 kinase regulatory subunit polypeptide 2; Phosphatidylinositol 3 kinase; regulatory subunit; polypeptide 2 (p85 beta); Phosphatidylinositol 3-kinase 85 kDa regulatory subunit beta; phosphatidylinositol 3-kinase; Phosphatidylinositol 3-kinase regulatory beta subunit; Phosphatidylinositol 3-kinase regulatory subunit beta; Phosphoinositide 3 kinase regulatory subunit 2 (beta); Phosphoinositide 3 kinase regulatory subunit 2; Phosphoinositide 3 kinase regulatory subunit polypeptide 2 (p85 beta); Phosphoinositide 3 kinase regulatory subunit polypeptide 2; Phosphoinositide 3 kinase; regulatory subunit 2 (beta); Phosphoinositide 3 kinase; regulatory subunit 2 (p85 beta); PI3 kinase p85 beta subunit; PI3 kinase p85 subunit beta; PI3-kinase regulatory subunit beta; PI3-kinase subunit p85-beta; PI3K; PI3K regulatory subunit beta; PIK3R 2; PIK3R2; polypeptide 2 (p85 beta); PtdIns 3 kinase p85 beta; PtdIns-3-kinase p85-beta; PtdIns-3-kinase regulatory subunit beta; PtdIns-3-kinase regulatory subunit p85-beta.
Gene ID	5296
Gene Name	PIK3R2
SwissProt ID	O00459
Host	Rabbit
Reactivity	Human, Rat
Molecular Weight	Calculated MW: 82 kDa; Observed MW: 82 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-13B9G6
Form	Liquid
Concentration	See label

Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human PI 3 Kinase p85 beta
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P, ICC/IF, FC, IP
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/50 FC: 1/50-1/100
Research Field	Signal Transduction
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

