

# **Anti-PKN1 Rabbit Monoclonal Antibody**

## Introduction

The protein encoded by this gene belongs to the protein kinase C superfamily. This kinase is activated by Rho family of small G proteins and may mediate the Rho-dependent signaling pathway. This kinase can be activated by phospholipids and by limited proteolysis. The 3-phosphoinositide dependent protein kinase-1 (PDPK1/PDK1) is reported to phosphorylate this kinase, which may mediate insulin signals to the actin cytoskeleton. The proteolytic activation of this kinase by caspase-3 or related proteases during apoptosis suggests its role in signal transduction related to apoptosis. Alternatively spliced transcript variants encoding distinct isoforms have been observed.

## Product parameters

Alternative Names	DBK; PKN; PAK1; PAK-1; PRKCL1; PKN-ALPHA
Gene ID	5585
Ge <mark>ne N</mark> ame	PKN1 ()
SwissProt ID	Q16512
Host	Rabbit
Reactivity	Rat
Molecular Weight	Calculated MW: 104 kDa; Observed MW: 120 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-13D3A4
Form	Liquid
Con <mark>cent</mark> ration	See label
Carrier	Carrier Not Free
Immunogen	A synthetic peptide of human PKN
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB

Dilution Ratio	WB: 1/500-1/1000
Research Field	Signal Transduction
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note A P	Please avoid freeze-thaw cycles.

### Protocol

Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

### Note

1. This product is for scientific research use only.

















