

## Anti-PPAR gamma Rabbit Monoclonal Antibody

### Introduction

Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the receptor binds to a promoter element in the gene for acyl-CoA oxidase and activates its transcription. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids. Key regulator of adipocyte differentiation and glucose homeostasis.

### Product parameters

Alternative Names	CIMT1; GLM1; NR1C3; PPAR gamma; PARG; PPARG1; PPARG2;
Gene ID	5468
Gene Name	PPARG
SwissProt ID	P37231
Host	Rabbit
Reactivity	Human
Molecular Weight	Calculated MW: 58 kDa; Observed MW: 58 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-9C5D1
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Peptide
Purification	Affinity Purified
Buffer System	Liquid in 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Application	WB, ICC/IF, FC
Dilution Ratio	WB: 1/500-1/1000 IF: 1/50-1/200 FC: 1/50-1/100
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

## Protocol



Configure the product according to the application range and recommended dilution ratio.

**\*Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.



**APExBIO Technology**

**[www.apexbt.com](http://www.apexbt.com)**

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: [info@apexbt.com](mailto:info@apexbt.com)

