

# **Anti-PSMA Rabbit Monoclonal Antibody**

#### Introduction

PSMA has both folate hydrolase and N-acetylated-alpha-linked-acidic dipeptidase (NAALADase) activity. Has a preference for tri-alpha-glutamate peptides. In the intestine, required for the uptake of folate. In the brain, modulates excitatory neurotransmission through the hydrolysis of the neuropeptide, N-aceylaspartylglutamate (NAAG), thereby releasing glutamate. Isoform PSM-4 and isoform PSM-5 would appear to be physiologically irrelevant. Involved in prostate tumor progression. Also exhibits a dipeptidyl-peptidase IV type activity. In vitro, cleaves Gly-Pro-AMC.

## Product parameters

Alternative Names	FGCP; FOLH 1; GCP 2; GCPII; mGCP; NAALADase I; PSM; PSMA	
Gene ID	2346	
Gene Name	FOLH1	
SwissProt ID	Q04609	
Host Achieve Perfec	Rabbit	
Reactivity	Human	
Molecular Weight	Calculated MW: 84 kDa; Observed MW: 110 kDa	
Conjugation	Unconjugated	
Ex	-	
Em	-	
Modification	Unmodified	
Clonality	IgG	
Isotype	Monoclonal Antibody	
Clonality No.	AP-1G2A2	
Form	Liquid	
Concentration	See label	
Carrier	Carrier Free	
Immunogen	A synthesized peptide derived from human PSMA	
Purification	Affinity Chromatography	
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	
Application	WB, IHC-P, IP	
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/50	

Research Field	Signal Transduction
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.





#### Protocol

Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

### Note

1. This product is for scientific research use only.

















