

Anti-RNA Helicase A Rabbit Monoclonal Antibody

Unwinds double-stranded DNA and RNA in a 3' to 5' direction. Alteration of secondary structure may subsequently influence interactions with proteins or other nucleic acids. Functions as a transcriptional activator. Component of the CRD-mediated complex that promotes MYC mRNA stability.

Product parameters

Introduction

| Alternative Names | LKP; RHA; DDX9; NDH2; NDHII |
|------------------------------|---|
| Gene ID | 1660 |
| Gene Name | DHX9 |
| SwissProt ID | Q08211 |
| Host | Rabbit |
| Reactivity | Rat |
| Molecular Weight | Calculated MW: 141 kDa; Observed MW: 141 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-20B3E1 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Not Free |
| Immunogen | A synthetic peptide of human RNA Helicase A |
| Purification | Affinity Purified |
| Buff <mark>er Sy</mark> stem | 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA. |
| Application | WB |
| Dilution Ratio | WB: 1/500-1/1000 |
| Research Field | Epigenetics and Nuclear Signaling |
| Product Categories | Primary antibody |
| Shipping | Blue ice |

| Storage | -20°C |
|-----------------|----------------------------------|
| Expiration Date | 12 months |
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





