

## **Anti-RON Rabbit Monoclonal Antibody**

#### Introduction

Ron is initially synthesized in the cells as a single-chain, pro-Ron precursor that is cleaved into the two active chains. RON signaling activates the wound healing response by promoting epithelial cell migration, proliferation as well as survival at the wound site. Plays also a role in the innate immune response by regulating the migration

### Product parameters

and phagocytic activity of macrophages.

Alternative Names	MST1R; PTK8; RON; Macrophage-stimulating protein receptor; MSP receptor; CDw136; Protein-tyros kinase 8; p185-Ron; CD136
Gene ID	4486
Gene Name	MST1R
SwissProt ID	Q04912
Host	Rabbit
Reactivity	Human Achieve Perfection, Explore the Unknown
Molecular Weight	Calculated MW: 152 kDa; Observed MW: 185 kDa
Conjugation	Unconjugated
Ex	-
Em	
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-1B9G7
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human RON
Purification Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P, ICC/IF, FC, IP
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/30 FC: 1/50-1/100
Research Field	Cell Biology

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

# | Protocol



Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

#### Note

1. This product is for scientific research use only.





