

## Anti-RPA32 Rabbit Monoclonal Antibody

### Introduction

As part of the heterotrimeric replication protein A complex (RPA/RP-A), binds and stabilizes single-stranded DNA intermediates, that form during DNA replication or upon DNA stress. It prevents their reannealing and in parallel, recruits and activates different proteins and complexes involved in DNA metabolism. Thereby, it plays an essential role both in DNA replication and the cellular response to DNA damage. In the cellular response to DNA damage, the RPA complex controls DNA repair and DNA damage checkpoint activation.

### Product parameters

Alternative Names	60S acidic ribosomal protein P1; AA409079; AI325195; AU020965; HSSB; ik:tdsubc_2g1; M(2)21C; MGC137236; OTTHUMP00000004008; p32; p34; RCJMB04_6d17 replication protein A2; 32kDa; REPA 2; REPA1; REPA2; Replication factor A protein 2; Replication protein A 32 kDa subunit; Replication protein A 32kDa subunit; Replication protein A 34 kDa subunit; Replication protein A; replication protein A1 (70kD); Replication Protein A2 (32kDa); Replication protein A2 32kD; Replication protein A2 32kDa; Replication protein A2; Replication protein A2; 32kDa; RF A; RF-A protein 2; Rf-A2; RFA; RFA2_HUMAN; RP A; RP-A p32; RP-A p34; RP21C; RPA 2; RPA 32; RPA; RPA2; RPA32; RPA34; RPA70; RpLP1; RpP2; xx:tdsubc_2g1; zgc:109822.
Gene ID	6118
Gene Name	RPA2
SwissProt ID	P15927
Host	Rabbit
Reactivity	Human, Rat
Molecular Weight	Calculated MW: 29 kDa; Observed MW: 29 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-6E7C12
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	A synthetic peptide of human RPA32/RPA2

Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB, IHC-P
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

## Protocol

Configure the product according to the application range and recommended dilution ratio.

**\*Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.





**APExBIO Technology**

**[www.apexbt.com](http://www.apexbt.com)**

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: [info@apexbt.com](mailto:info@apexbt.com)

