

## Anti-S100A1 Rabbit Monoclonal Antibody

### Introduction

Weakly binds calcium but binds zinc very tightly-distinct binding sites with different affinities exist for both ions on each monomer. Physiological concentrations of potassium ion antagonize the binding of both divalent cations, especially affecting high-affinity calcium-binding sites.

### Product parameters

Alternative Names	S100A1; S100A; Protein S100-A1; S-100 protein alpha chain; S-100 protein subunit alpha; S100 calcium-binding protein A1
Gene ID	6271
Gene Name	S100A1
SwissProt ID	P23297
Host	Rabbit
Reactivity	Human, Mouse
Molecular Weight	Calculated MW: 11 kDa; Observed MW: 11 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-1C1H8
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human S100
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P, FC, IP
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/50 FC: 1/50-1/100
Research Field	Cell Biology
Product Categories	Primary antibody
Shipping	Blue ice

Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

## Protocol

Configure the product according to the application range and recommended dilution ratio.

**\*Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.



**APExBIO Technology**

**[www.apexbt.com](http://www.apexbt.com)**

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: [info@apexbt.com](mailto:info@apexbt.com)

