

Anti-SETD7 Rabbit Monoclonal Antibody

Introduction

Histone methyltransferase that specifically monomethylates 'Lys-4' of histone H3. H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation. Plays a central role in the transcriptional activation of genes such as collagenase or insulin. Recruited by IPF1/PDX-1 to the insulin promoter, leading to activate transcription.

Product parameters

Alternative Names	Histone H3-K4 methyltransferase SETD7; Lysine N-methyltransferase 7; SET domain-containing protein 7; SET7; SET9
Gene ID	80854
Gene Name	SETD7
SwissProt ID	Q8WTS6
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 41 kDa; Observed MW: 41,50 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-6G10D10
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Peptide
Purification	Affinity Purified
Buffer System	Liquid in 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Application	WB, IP
Dilution Ratio	WB: 1/500-1/1000 IP: 1/20
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol



Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

