


events in T-lymphocytes are different from those in B-cells.

Anti-SLAMF1 Rabbit Monoclonal Antibody

Introduction



Product parameters

Alternative Names	SLAMF1; SLAM; Signaling lymphocytic activation molecule; CDw150; IPO-3; CD150
Gene ID	6504
Gene Name	SLAMF1
SwissProt ID	Q13291
Host	Rabbit
Reactivity	Human
Molecular Weight	Calculated MW: 37 kDa; Observed MW: 85 kDa
Conjugation	Unconjugated A P E P
Ex Achieve Perfect	Achieve Perfection, Explore the Unknown
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-1F2F12
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human SLAMF1
Purification	Affinity Purified
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, FC, IP
Dilution Ratio	WB: 1/500-1/1000 IP: 1/50 FC: 1/50-1/100
Research Field	-
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
	.i.

Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





