

Anti-STAT2 Rabbit Monoclonal Antibody

Introduction

STAT2 (113-kDa), originally purified from the nuclei of alpha-interferon-treated cells, is critical to the transcriptional responses induced by type I interferons, IFN-alpha/beta. Stat2 is rapidly activated by phosphorylation at Tyr690 in response to stimulation by IFN-alpha/beta via associations with receptor-bound Jak kinases

Product parameters

Alternative Names	Homo sapiens interferon alpha induced transcriptional activator; ISGF 3; P113; signal transducer an activator of transcription 2 113kD; STAT113; Stat2
Gene ID	6773
Gene Name	STAT2
SwissProt ID	P52630
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 98 kDa; Observed MW: 98 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-12G5C9
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Im <mark>mu</mark> nogen	A synthesized peptide derived from human STAT2
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P, ICC/IF
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200
Research Field	Signal Transduction
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol



Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





