

## **Anti-STAT5 Rabbit Monoclonal Antibody**

#### Introduction



Carries out a dual function: signal transduction and activation of transcription. Mediates cellular responses to the cytokine KITLG/SCF and other growth factors. May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4. Binds to the GAS element and activates PRL-induced transcription. Regulates the expression of milk proteins during lactation.

### Product parameters

Alternative Names	MGF; STAT5
Gene ID	20850
Gene Name	Stat5A
SwissProt ID	P42230
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 91 kDa; Observed MW: 91 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	lgG
Isotype	Monoclonal Antibody
Clonality No.	AP-7C1F7
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Recombinant protein of mouse STAT5A
Pu <mark>rifica</mark> tion	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB, ICC/IF, IP
Dilution Ratio	WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/20
Research Field	Signal Transduction

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

# | Protocol



Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

#### Note

1. This product is for scientific research use only.





