

Anti-STAT6 Rabbit Monoclonal Antibody

STAT6 transcription factor of the STAT family. Plays a central role in IL4-mediated biological responses. Induces the expression of BCL2L1/BCL-X(L), which is responsible for the anti-apoptotic activity of IL4.

Product parameters

Introduction

| Alternative Names | STAT6; Signal transducer and activator of transcription 6; IL-4 Stat |
|--------------------|--|
| Gene ID | 6778 |
| Gene Name | STAT6 |
| SwissProt ID | P42226 |
| Host | Rabbit |
| Reactivity | Human, Mouse |
| Molecular Weight | Calculated MW: 94 kDa |
| Conjugation | Unconjugated |
| Ex Achieve Perfect | Explore the Unknown |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-19G4F11 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Free |
| Immunogen | A synthesized peptide derived from human STAT6 |
| Purification | Affinity Chromatography |
| Buffer System | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol |
| Application | WB, IHC-P, ICC/IF, IP |
| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/50 |
| Research Field | Signal Transduction |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |

| Expiration Date | 12 months |
|-----------------|----------------------------------|
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





