

Anti-Sulfotransferase 2A1 Rabbit Monoclonal Antibody

Introduction

Catalyzes the sulfation of steroids and bile acids in the liver and adrenal glands.

Product parameters

| | |
|--------------------|--|
| Alternative Names | HST; ST2; STD; hSTa; DHEAS; ST2A1; ST2A3; DHEA-ST |
| Gene ID | 6822 |
| Gene Name | SULT2A1 |
| SwissProt ID | Q06520 |
| Host | Rabbit |
| Reactivity | Human, Mouse, Rat |
| Molecular Weight | Calculated MW: 34 kDa; Observed MW: 34 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-7C4A3 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Not Free |
| Immunogen | Peptide |
| Purification | Affinity Purified |
| Buffer System | Liquid in 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. |
| Application | WB, IHC-F, IHC-P, ICC/IF, FC |
| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 FC: 1/50-1/100 |
| Research Field | Signal Transduction |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |
| Expiration Date | 12 months |

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

