

## Anti-Toll-Like Receptor 7 Rabbit Monoclonal Antibody

### Introduction

Key component of innate and adaptive immunity. TLRs (Toll-like receptors) control host immune response against pathogens through recognition of molecular patterns specific of microorganisms. TLR7 is a nucleotide-sensing TLR which is activated by single-stranded RNA. Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response.

### Product parameters

Alternative Names	Toll-Like Receptor 7; PRO285; TLR 7; Tlr7; Toll like receptor 7; UNQ248
Gene ID	51284
Gene Name	TLR7
SwissProt ID	Q9NYK1
Host	Rabbit
Reactivity	Human
Molecular Weight	Calculated MW: 121 kDa; Observed MW: 140 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-9E4C12
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human TLR7
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100
Research Field	Immunology
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

## Protocol



Configure the product according to the application range and recommended dilution ratio.

**\*Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.



**APExBIO Technology**

**[www.apexbt.com](http://www.apexbt.com)**

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: [info@apexbt.com](mailto:info@apexbt.com)

