

Anti-Transcription Factor AP 2 gamma Rabbit Monoclonal Antibody

Introduction

The protein encoded by this gene is a sequence-specific DNA-binding transcription factor involved in the activation of several developmental genes. The encoded protein can act as either a homodimer or heterodimer with other family members and is induced during retinoic acid-mediated differentiation. It plays a role in the development of the eyes, face, body wall, limbs, and neural tube.

Product parameters

Alternative Names	ERF1; TFAP2G; hAP-2g; AP2-GAMMA
Gene ID	7022
Gene Name	TFAP2C
SwissProt ID	Q92754
Host	Rabbit
Reactivity	Human
Molecular Weight	-
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-17A5H5
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Recombinant protein of human AP2 gamma
Purification	Affinity Purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	IHC-P
Dilution Ratio	IHC: 1/100-1/200
Research Field	Cell Biology
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol



Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

