

Anti-Transcription Initiation Factor IIB (2F6) Mouse Monoclonal Antibody

Introduction

General factor that plays a major role in the activation of eukaryotic genes transcribed by RNA polymerase II.

Product parameters

Alternative Names	General transcription factor IIB; General transcription factor TFIIB; gtf2b; RNA polymerase II transcription factor IIB; S300 II; S300-II; TF IIB; TF2B; TF2B_HUMAN; TFIIB; Transcription initiation factor IIB.
Gene ID	2959
Gene Name	GTF2B
SwissProt ID	Q00403
Host	Mouse
Reactivity	Human, Mouse, Monkey
Molecular Weight	Calculated MW: 35 kDa; Observed MW: 35 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG2b
Isotype	Monoclonal Antibody
Clonality No.	AP-17C6D5
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Purified recombinant human TFIIB protein fragments expressed in E.coli.
Purification	Affinity Purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	WB
Dilution Ratio	WB: 1/500-1/1000
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

