

## Anti-TRPM8 Rabbit Monoclonal Antibody

Activated by icilin, eucalyptol, menthol, cold and modulation of intracellular pH. Involved in menthol sensation. Permeable for monovalent cations sodium, potassium, and cesium and divalent cation calcium.

## Product parameters

Introduction

Alternative Names	LTrpC6; MGC2849; Transient receptor potential p8; Trp p8; Trpm8; TRPP8
Gene ID	79054
Gene Name	TRPM8
SwissProt ID	Q7Z2W7
Host	Rabbit
Reactivity	Human, Mouse
Molecular Weight	Calculated MW: 128 kDa; Observed MW: 128 kDa
Conjugation	Unconjugated
Ex Actueve Perfect	ch, Explore the Unknown
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-11G2A1
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human TRPM8
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol
Application	WB, ICC/IF
Dilution Ratio	WB: 1/500-1/1000 IF: 1/50-1/200
Research Field	Signal Transduction
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C

Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

## Protocol

Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.





