

Anti-Tuberin Rabbit Monoclonal Antibody

Introduction



Tuberin is a product of the TSC2 tumor suppressor gene and an important regulator of cell proliferation and tumor development. Mutations in either TSC2 or the related TSC1 (hamartin) gene cause tuberous sclerosis complex (TSC), an autosomal dominant disorder characterized by development of multiple, widespread non-malignant tumors. Tuberin is directly phosphorylated at Thr1462 by Akt/PKB. Phosphorylation at Thr1462 and Tyr1571 regulates tuberin-hamartin complexes and tuberin activity.

Product parameters

Alternative Names	TSC2; Tuberous sclerosis 2 homolog protein; Tuberous sclerosis 2 protein; Tuberin; TSC4	
Gene ID	7249	
Gene Name	TSC2	
SwissProt ID	P49815	
Host	Rabbit	
Reactivity	Human Addition Explor	
Molecular Weight	Calculated MW: 201 kDa; Observed MW: 201 kDa	
Conjugation	Unconjugated	
Ex	-	
Em	-	
Modification	Unmodified	
Clonality	IgG	
Isotype	Monoclonal Antibody	
Clonality No.	AP-5B5H11	
Form	Liquid	
Concentration	See label	
Carrier	Carrier Not Free	
Immunogen	Recombinant protein of human Tuberin	
Purification	Affinity Purified	
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.	
Application	WB, IP	
Dilution Ratio	WB: 1/500-1/1000 IP: 1/20	
Research Field	Cell Biology	

Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.





Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





