

Introduction

Anti-UFD1 Rabbit Monoclonal Antibody

Essential component of the ubiquitin-dependent proteolytic pathway which degrades ubiquitin fusion proteins. The ternary complex containing UFD1, VCP and NPLOC4 binds ubiquitinated proteins and is necessary for the export of misfolded proteins from the ER to the cytoplasm, where they are degraded by the proteasome. The NPLOC4-UFD1-VCP complex regulates spindle disassembly at the end of mitosis and is necessary for the formation of a closed nuclear envelope. It may be involved in the development of some ectoderm-derived structures . Acts as a negative regulator of type I interferon production via the complex formed with VCP and NPLOC4, which binds to DDX58/RIG-I and recruits RNF125 to promote ubiquitination and degradation of DDX58/RIG-I (PubMed:26471729).

Product parameters

Alternative Names	UFD1L
Gene ID	
Gene Name	UFD1 Acheve Perfection, Explore the Unknown
SwissProt ID	Q92890
Host	Rabbit
Reactivity	Human
Molecular Weight	Calculated MW: 35 kDa; Observed MW: 40 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-4C6F4
Form	
Concentration	See label
Carrier	Carrier Not Free
Immunogen	A synthetic peptide of human UFD1L
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.

Application	WB	
Dilution Ratio	WB: 1/500-1/1000	
Research Field	Cell Biology	
Product Categories	Primary antibody	
Shipping	Blue ice	
Storage	-20°C	
Expiration Date		
Note	Please avoid freeze-thaw cycles.	

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



















