

Anti-USP22 Rabbit Monoclonal Antibody

Introduction

Histone deubiquitinating component of the transcription regulatory histone acetylation (HAT) complex SAGA. Catalyzes the deubiquitination of both histones H2A and H2B, thereby acting as a coactivator.

Product parameters

| | |
|--------------------|---|
| Alternative Names | Ubiquitin carboxyl terminal hydrolase 22; Usp22; USP3L |
| Gene ID | 23326 |
| Gene Name | USP22 |
| SwissProt ID | Q9UPT9 |
| Host | Rabbit |
| Reactivity | Human, Mouse, Rat |
| Molecular Weight | Calculated MW: 60 kDa; Observed MW: 60 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-10C2F1 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Free |
| Immunogen | A synthesized peptide derived from human USP22 |
| Purification | Affinity Chromatography |
| Buffer System | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Application | WB, ICC/IF, IP |
| Dilution Ratio | WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/50 |
| Research Field | Epigenetics and Nuclear Signaling |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |

| | |
|-----------------|----------------------------------|
| Expiration Date | 12 months |
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APEx BIO Technology
www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

