

Anti-C-Myc Tag (9F6) Mouse Monoclonal Antibody

Introduction

Myc tag is a polypeptide protein tag derived from the c-myc gene product that can be added to a protein using recombinant DNA technology. It can be used for affinity chromatography, and then used to separate recombinant, overexpressed protein from wild type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits.

Product parameters

Alternative Names	c-Myc; Myc tag; Myc protein; Myc epitope tag; Myc2
Gene ID	-
Gene Name	-
SwissProt ID	-
Host	Mouse
Reactivity	Species-independent
Molecular Weight	-
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG1
Isotype	Monoclonal Antibody
Clonality No.	AP-3A3H6
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Synthetic peptide EQKLISEEDL coupled to KLH.
Purification	Affinity Purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	WB
Dilution Ratio	WB: 1/1000-1/10000
Research Field	Tags & Cell Markers
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol



Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

