

Anti-His Tag (2A2) Mouse Monoclonal Antibody

Introduction

A polyhistidine-tag is an amino acid motif in proteins that consists of at least five histidine (His) residues, often at the N- or C-terminus of the protein. It is also known as hexa histidine-tag, 6xHis-tag, and by the trademarked name His-tag. Polyhistidine-tags are often used for affinity purification of polyhistidine-tagged recombinant proteins expressed in Escherichia coli and other prokaryotic expression systems.

Product parameters

Alternative Names	6 His epitope tag; Hexa His tag; HHHHHH epitope tag; HHHHHH tag; His tag
Gene ID	-
Gene Name	-
SwissProt ID	-
Host	Mouse
Reactivity	Species-independent
Molecular Weight	-
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG1
Isotype	Monoclonal Antibody
Clonality No.	AP-17E4F6
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Synthetic peptide: HHHHHH conjugated to KLH.
Purification	Affinity Purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	WB, ICC/IF, IP
Dilution Ratio	WB: 1/1000-1/10000 IF: 1/100-1/500 IP: 1/200
Research Field	Tags & Cell Markers
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol



Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APEx BIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

