

Recombinant Mouse KGF-2/FGF-10

Information

Gene ID	14165
Accession #	O35565
Alternate Names	
Source	<i>Escherichia coli</i> .
M.Wt	Approximately 19.5 kDa, a single non-glycosylated polypeptide chain containing 173 amino acids.
AA Sequence	QALGQDMVSQ EATNCSSSSS SFSSPSSAGR HVRSYNHLQG DVRWRRLFSF TKYFLTIEKN GKVSGTKNED CPYSVLEITS VEIGVVAVKA INSNYLAMN KKGKLYGSKE FNNDCKLKER IEENGYNTYA SFNWQHNGRQ MYVALNGKGA PRRGQKTRRK NTSAHFLPMT IQT
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. - 12 months from date of receipt, -20 to -70 °C as supplied. - 1 month, 2 to 8 °C under sterile conditions after reconstitution. - 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 2 × PBS, 600 mM NaCl, pH 7.4, 1 mM mercaptoethanol.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by thymidine uptake assay using FGF-receptors transfected BaF3 cells is less than 0.5 ng/ml, corresponding to a specific activity of > 2.0 × 10 ⁶ IU/mg.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	5 µg	100 µg	500 µg
Recombinant Mouse KGF-2/FGF-10	5 µg	100 µg	500 µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.

Quality Control

Purity	> 95 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 1 EU/μg of rMuKGF-2/FGF-10 as determined by LAL method.

Description

Fibroblast growth factor 10 belongs to the fibroblast growth factor (FGF) family, which is involved in a variety of biological processes such as embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. Like most other FGF family members, FGF-10 also has a heparin-binding domain and plays an important role in the regulation of embryonic development, cell proliferation and cell differentiation. In addition, FGF-10 may take parts in wound healing and is required for normal branching morphogenesis. Recombinant murine FGF-10 contains a 173 amino acids and it shares 94 % and 100 % a.a. sequence identity with human and rat FGF-10.

Reference

1. Emoto H, Tagashira S, Mattei MG, et al. 1997. J Biol Chem. 272:23191-4.
2. Tagashira S, Harada H, Katsumata T, et al. 1997. Gene. 197:399-404.
3. Carninci P, Kasukawa T, Katayama S, et al. 2005. Science. 309:1559-63.
4. Igarashi M, Finch PW, Aaronson SA. 1998. J Biol Chem. 273:13230-5.

APEx BIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com