

Recombinant Human Neuregulin 1-alpha EGF Domain

Information

Gene ID	3084
Accession #	Q02297
Alternate Names	
Source	Escherichia coli.
M.Wt	Approximately 7.4 kDa, a single non-glycosylated polypeptide chain containing 65 amino acids.
AA Sequence	SHLVKCAEKE KTFCVNGGEC FMVKDLSNPS RYLCKCQPGF TGARCTENVP MKVQNQEKAE ELYQK
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution
Formulation	Lyophilized from a 0.2 μm filtered solution in 20 mM PB, pH 6.0, 150 mM NaCl.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at \leq -20 °C. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The ED as determined by a cell proliferation assay using serum free human MCF-7 cells is less than 40 ng/ml, corresponding to a specific activity of $> 2.5 \times 10$ IU/mg.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

DOLE R		Perfect	
Components Contract State	10µg	100µg	500µg
Recombinant Human Neuregulin 1-alpha EGF Domain	10µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

I,	Quality Control	19. 	
	Purity	> 97 % by SDS-PAGE and HPLC analyses.	
	Endotoxin	Less than 0.1 EU/µg of rHuNRG1- α as determined by LAL method.	

Description

Reference



APENER BIO





