

Recombinant Human Basic Fibroblast Growth Factor GMP

Information

Gene ID	2247
Accession #	P09038
Alternate Names	FGF-2, HBGF-2
Source	Escherichia coli.
M.Wt	Approximately 16.5 kDa, a single non-glycosylated polypeptide chain containing 147 amino acids.
AA Sequence	MPALPEDGGS GAFPPGHFKD PKRLYCKNGG FFLRIHPDGR VDGVREKSDP HIKLQLQAE E RGVVSIKGVC ANRYLAMKED GRLLASKCVT DECFFFERLE SNNYNTYRSR KYTSWYVALK RTGQYKLGSK TGPQGKAILF LPMSAKS
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles A minimum of 12 months when stored at $\leq -20^{\circ}\text{C}$ as supplied Refer to lot specific COA for the Use by Date 1 month, 2 to 8°C under sterile conditions after reconstitution 3 months, -20 to -70°C under sterile conditions after reconstitution
Formulation	Lyophilized from a $0.2\ \mu\text{m}$ filtered concentrated solution in 20 mM Tris-HCl, pH 7.6, with 150mM NaCl.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of less than 0.3 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $\leq -20^{\circ}\text{C}$. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The ED as determined by a cell proliferation assay using murine balb/c 3T3 cells is less than 0.05 ng/ml, corresponding to a specific activity of $> 2.0 \times 10\ \text{IU/mg}$.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	5 μg	100 μg	
Recombinant Human Basic Fibroblast Growth Factor GMP	5 μg	100 μg	

Use a manual defrost freezer and avoid repeated freeze-thaw cycles
 A minimum of 12 months when stored at $\leq -20^{\circ}\text{C}$ as supplied
 Refer to lot specific COA for the Use by Date
 1 month, 2 to 8°C under sterile conditions after reconstitution
 3 months, -20 to -70°C under sterile conditions after reconstitution

Quality Control

Purity	> 98 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 0.01 EU/ μg of rHubFGF GMP as determined by LAL method.

Description

Human bFGF, encoded by the FGF2 gene, is a member of the fibroblast growth factor (FGF) family. Fibroblast growth factor was found in pituitary extracts in 1973 and then tested in a bioassay that caused fibroblasts to proliferate. After further fractionating the extract using acidic and basic pH, two different forms have isolated that named "acidic fibroblast growth factor" (FGF-1) and "basic fibroblast growth factor" (FGF-2). Human bFGF shares 54 % amino acid sequence identity with aFGF. Affinity between bFGF and its receptors can be increased by heparin or heparan sulfate proteoglycan. bFGF plays an important role in the regulation of cell survival, cell division, angiogenesis, cell differentiation and cell migration. bFGF are also involved in a variety of biological processes, including embryonic development, morphogenesis, tissue repair, tumor growth and invasion. Additionally, bFGF is frequently used for a critical component of cell culture medium, e.g., human embryonic stem cell culture medium, serum-free culture systems.

Reference

1. Armelin HA. 1973. Proc Natl Acad Sci U S A. 70:2702-6
2. Gospodarowicz D. 1974. Nature. 249:123-7
3. Eswarakumar VP, Lax I, Schlessinger J. 2005. Cytokine Growth Factor Rev. 16:139-49
4. Ornitz DM, Xu J, Colvin JS, et al. 1996. J Biol Chem. 271:15292-7
5. Landriscina M, Bagala C, Mandinova A, et al. 2001. J Biol Chem. 276:25549-57
6. Fernandez IS, Cuevas P, Angulo J, et al. 2010. J Biol Chem. 285:11714-29
7. Liu Y, Song Z, Zhao Y, et al. 2006. Biochem Biophys Res Commun. 346:131-9.

APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com