

Recombinant Human Keratinocyte Growth Factor-1/FGF-7 GMP

Information

Gene ID	2252
Accession #	P21781
Alternate Names	
Source	Escherichia coli.
M.Wt	Approximately 18.9 kDa, a single, non-glycosylated polypeptide chain containing 163 amino acids.
AA Sequence	CNDMTPEQMA TNVNCSSPER HTRSYDMEG GDIRVRLFC RTQWYLRIK RGKVKGQTQEM KNNYNIMEIR TVAVGIVAIAK GVESEFYLAM NKEGKLYAKK ECNEDCNFKE LILENHNTY ASAKWTHNGG EMFVALNQKG IPVRGKKTCK EQKTAHFLPM AIT
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles A minimum of 12 months when stored at $\leq -20^{\circ}\text{C}$ as supplied Refer to lot specific COA for the Use by Date 1 month, 2 to 8°C under sterile conditions after reconstitution 3 months, -20 to -70°C under sterile conditions after reconstitution
Formulation	Lyophilized from a $0.2\ \mu\text{m}$ filtered solution in 20mM PB, pH 7.4, 500 mM NaCl, 0.02 % Tween-20.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $\leq -20^{\circ}\text{C}$. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The ED as determined by thymidine uptake assay using FGF-receptors transfected BaF3 cells is less than 10 ng/ml, corresponding to a specific activity of $> 1.0 \times 10\ \text{IU/mg}$.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	5 μg	100 μg	
Recombinant Human Keratinocyte Growth Factor-1/FGF-7 GMP	5 μg	100 μg	

Use a manual defrost freezer and avoid repeated freeze-thaw cycles
A minimum of 12 months when stored at $\leq -20^{\circ}\text{C}$ as supplied
Refer to lot specific COA for the Use by Date
1 month, 2 to 8°C under sterile conditions after reconstitution
3 months, -20 to -70°C under sterile conditions after reconstitution

Quality Control

Purity	> 98 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 0.01 EU/ μg of rHuKGF-1/FGF-7 GMP as determined by LAL method.

Description

Reference

APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com