

Recombinant Human VEGF Co-regulated Chemokine 1/CXCL17

Information

Gene ID	284340	
Accession #	Q6UXB2	
Alternate Names		
Source	Escherichia coli.	
M.Wt	Approximately 11.5 kDa, a single non-glycosylated polypeptide chain containing 98 amino acids.	
AA Sequence	SSLNPGVARG HRDRGQASRR WLQEGGQECE CKDWFLRAPR RKFMTVSGLP KKQCPCDHFK GNVKKTRHQR HHRKPNKHSR ACQQFLKQCQ LRSFALPL	
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.	
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution	
Formulation	Lyophilized from a 0.2 μ m filtered concentrated solution in PBS, pH 7.4, with 3 % Trehalose.	
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.	
Biological Activity	Fully biologically active when compared to standard. The ED as determined by its ability to induce VEGF expression using murine endothelial cells is less than 5.0 μ g/ml, corresponding to a specific activity of > 200 IU/mg.	
Shipping Condition	Gel pack.	
Handling	Centrifuge the vial prior to opening.	
Usage	For Research Use Only! Not to be used in humans.	

Components and Storage

Components	5µg	100µg	500µg
Recombinant Human VEGF Co-regulated Chemokine 1/CXCL17	5µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

Quality Control

	A Strong	akino.
Purity	> 96 % by SDS-PAGE and HPLC analyses.	Production and the state of the
Endotoxin	Less than 0.1 EU/μg of rHuVCC-1/CXCL17 a	s determined by LAL method.

Description

Reference





10





APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com