

Recombinant Human Hemofiltrate CC Chemokine-1/CCL14

Information

Gene ID	6358
Accession #	Q16627
Alternate Names	HCC-1/HCC-3, NCC-2, Small-inducible Cytokine A14
Source	Escherichia coli.
M.Wt	Approximately 8.4 kDa, a single non-glycosylated polypeptide chain containing 72 amino acids.
AA Sequence	TESSSRGPYH PSECCFTYTT YKIPRQRIMD YYETNSQCSK PGIVFITKRG HSVCTNPSDK WVQDYIKDMK EN
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 100 mM NaCl.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human monocytes is in a concentration of 5.0-20 ng/ml.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	10µg	100µg	500µg
Recombinant Human Hemofiltrate CC Chemokine-1/CCL14	10µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

Quality Control

Purity	> 96 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 1 EU/μg of rHuHCC-1/CCL14 as determined by LAL method.

Description

Human CCL14 is belonging to the CC chemokine family. It has two isoforms, CCL14a (HCC-1) and CCL14b (HCC-3). The sequence of HCC-3 differs from HCC-1 as follow: 27-27 R→ QTGGKPKVVKIQLKLVG. CCL14 was first isolated from the hemofiltrate of human patients with chronic renal failure. CCL14 promotes chemotaxis of T lymphocytes, monocytes and eosinophils, and inhibits infection of M-tropic human immunodeficiency virus type 1 and is a ligand for CCR1, CCR3 and CCR5. CCL14 is expressed constitutively in several normal tissues: spleen, liver, skeletal and heart muscle, gut, and bone marrow, present at high concentrations in plasma.

Reference

1. Schulz-Knappe P, Magert HJ, Dewald B, et al. 1996. J Exp Med. 183:295-9
2. Detheux M, Standker L, Vakili J, et al. 2000. J Exp Med. 192:1501-8
3. Pardigol A, Forssmann U, Zucht HD, et al. 1998. Proc Natl Acad Sci U S A. 95:6308-13.

APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com