

Recombinant Human Macrophage Inflammatory Protein-5, 68a.a./CCL15

Information

Gene ID	6359
Accession #	Q16663
Alternate Names	
Source	Escherichia coli.
M.Wt	Approximately 7.4 kDa, a single non-glycosylated polypeptide chain containing 68 amino acids.
AA Sequence	SFHFAADCCT SYISQSIPCS LMKSYPFETSS ECSKPGVIFL TTKGRQVCAK PSGPGVQDCM KKLKPYSI
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution
Formulation	Lyophilized from a 0.2 μm filtered concentrated solution in 1 × PBS, pH 7.2.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration range of 1.0-10 ng/ml.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	5μg	100μg	500μg
Recombinant Human Macrophage Inflammatory Protein-5, 68a.a./CCL15	5μg	100μg	500μg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

Quality Control

Purity	> 98 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 0.1 EU/ μ g of rHuMIP-5, 68a.a./CCL15 as determined by LAL method.

Description

Reference

APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com