

Recombinant Human Eotaxin-2/CCL24

Information

Gene ID	6369
Accession #	O00175
Alternate Names	CK-beta-6, MPIF-2, Small-inducible Cytokine A24
Source	Escherichia coli.
M.Wt	Approximately 8.8 kDa, a single non-glycosylated polypeptide chain containing 78 amino acids.
AA Sequence	VVIPSPCCMF FVSKRIPENR VVS YQLSSRS TCLKAGVIFT TTKGQQFCGD PKQEWVQRYM KNLDAKQKKA SPRARAVA
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human peripheral blood eosinophils is in a concentration of 50-100 ng/ml.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	5µg	100µg	500µg
Recombinant Human Eotaxin-2/CCL24	5µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

Quality Control

Purity	> 97 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 1 EU/ μ g of rHuEotaxin-2/CCL24 as determined by LAL method.

Description

CCL24 is a secreted protein, encoded by CCL24 gene in humans located on Chr.7, and produced by activated monocytes and T lymphocytes. CCL24 has functions of chemotactic activity for resting T-lymphocytes and eosinophils, but none for monocytes and activated lymphocytes. The plasma levels of CCL24 and the aspirin-exacerbated respiratory disease (such as asthma) morbidity rate have positive correlation.

Reference

1. Owczarek W, Paplinska M, Targowski T, et al. 2010. Cytokine, 50: 181-5
2. Ben-Yehuda C, Bader R, Puxeddu I, et al. 2008. Exp Lung Res, 34: 467-79
3. Schaefer D, Meyer JE, Pods R, et al. 2006. Int Arch Allergy Immunol, 140: 205-14.

APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com