

Recombinant Rhesus Macaque Eotaxin/CCL11

Information

Gene ID					
Accession #	Q8MIT7				
Alternate Names					
Source	Escherichia coli.				
M.Wt	Approximately 8.4 kDa, a single non-glycosylated polypeptide chain containing 74 amino acids.				
AA Sequence	GPDSVATTCC FTLTNKKIPL QRLESYRRII SGKCPQKAVI FKTKLAKDIC ADPKKKWVQD SMKYLDRKSP TPKP				
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.				
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution				
Formulation	Lyophilized from a 0.2 μ m filtered concentrated solution in 2 $ imes$ PBS, pH 6.9.				
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.				
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human peripheral blood eosinophils is in a concentration range of 0.1-10.0 ng/ml.				
Shipping Condition	Gel pack.				
Handling	Centrifuge the vial prior to opening.				
Usage	For Research Use Only! Not to be used in humans.				
omponents and S	torage		A P La Barbara	Sungain	
Petilon Commence of College Commence of College Colleg		5ua	400.00	500ug	

Components and Storage

Components	5µg	100µg	500μց
Recombinant Rhesus Macaque Eotaxin/CCL11	5µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

Quality Control

	All the state of t	Nicho a.
Purity	> 98 % by SDS-PAGE and HPLC analyses.	P and the state of
Endotoxin	Less than 0.1 EU/µg of rRhEotaxin/CCL11 as	determined by LAL method.

Description

Reference





10





APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com