

## Recombinant Murine Cuteaneous T-cell Attracting Chemokine/CCL27

### Information

<b>Gene ID</b>	20301
<b>Accession #</b>	Q9Z1X0
<b>Alternate Names</b>	
<b>Source</b>	Escherichia coli.
<b>M.Wt</b>	Approximately 10.9 kDa, a single, non-glycosylated polypeptide chain containing 95 amino acids.
<b>AA Sequence</b>	LPLPSSTSCC TQLYRQPLPS RLLRRIVHME LQADGDCHL QAVVLHLARR SVCVHPQNRSLARWLERQGGRLQGTVPVSLN LVLQKKMYSN PQQQN
<b>Appearance</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Stability &amp; Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered concentrated solution in 2 × PBS, pH 7.4, with 5% trehalose.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
<b>Biological Activity</b>	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human peripheral blood lymphocytes is in a concentration range of 10-100 ng/ml.
<b>Shipping Condition</b>	Gel pack.
<b>Handling</b>	Centrifuge the vial prior to opening.
<b>Usage</b>	For Research Use Only! Not to be used in humans.

### Components and Storage

Components	5µg	100µg	500µg
Recombinant Murine Cuteaneous T-cell Attracting Chemokine/CCL27	5µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

## Quality Control

Purity	> 98 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 1 EU/μg of rMuCTACK/CCL27 as determined by LAL method.

## Description

## Reference

**APExBIO Technology**

**[www.apexbt.com](http://www.apexbt.com)**

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: [info@apexbt.com](mailto:info@apexbt.com)