

Recombinant Rat LPS-induced CXC Chemokine/CXCL5

Information

Gene ID	60665
Accession #	P97885
Alternate Names	Cytokine LIX, Small-inducible Cytokine B5
Source	Escherichia coli.
M.Wt	Approximately 10.1 kDa, a single, non-glycosylated polypeptide chain containing 93 amino acids.
AA Sequence	APFSAMVATE LRCVCLTLAP RINPKMIANL EVIPAGPHCP KVEVIAKLKN QKDNVCLDPQ APLIKKVIQK ILGSENKGTK RNALALVRSR STQ
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 2 × PBS, pH 7.4.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human peripheral blood neutrophils is in a concentration of 10-100 ng/ml.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	5µg	100µg	500µg
Recombinant Rat LPS-induced CXC Chemokine/CXCL5	5µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

Quality Control

Purity	> 97 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 1 EU/ μ g of rRtLIX/CXCL5 as determined by LAL method.

Description

CXCL5 is a small cytokine belonging to the CXC chemokine family that is also known as epithelial-derived neutrophil-activating peptide 78 (ENA-78). It is produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor-alpha. The expression of CXCL5 has also been observed in eosinophils, and can be inhibited with the type II interferon IFN- γ . Recombinant rat CXCL5 contains 93 amino acids which is a single, non-glycosylated polypeptide chain. Rat CXCL5 shares 57 % amino acid sequence identity with human CXCL5.

Reference

1. Chang MS, McNinch J, Basu R, et al. 1994. J Biol Chem. 269:25277-82
2. O'Donovan N, Galvin M, Morgan JG. 1999. Cytogenet Cell Genet. 84:39-42
3. Persson T, Monsef N, Andersson P, et al. 2003. Clin Exp Allergy. 33:531-7.

APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com