

# Recombinant Rat Breast and Kidney-expressed Chemokine/CXCL14

### Information

Gene ID	306748	
Accession #	Q8K453	
Alternate Names		
Source	Escherichia coli.	
M.Wt	Approximately 9.4 kDa, a single non-glycosylated polypeptide chain containing 77 amino acids.	
AA Sequence	SKCKCSRKGP KIRYSDVKKL EMKPKYPHCE EKMVIITTKS MSRYRGQEHC LHPKLQSTKR FIKWYNAWNE KRRVYEE	
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.	
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution	
Formulation	Lyophilized from a 0.2 $\mu$ m filtered concentrated solution in 30 % Acetonitrile an 0.1 % TFA.	
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.	
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human monocytes is in a concentration range of 1.0-10 ng/ml.	
Shipping Condition	Gel pack.	
Handling	Centrifuge the vial prior to opening.	
Usage	For Research Use Only! Not to be used in humans.	

## Components and Storage

Components	5µg	100µg	500µg
Recombinant Rat Breast and Kidney- expressed Chemokine/CXCL14	5µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

## Quality Control

	Akhou.	Nicho W
Purity	> 95 % by SDS-PAGE and HPLC analyses.	P. C. Land Control of the Control of
Endotoxin	Less than 0.1 EU/µg of rRtBRAK/CXCL14 as	determined by LAL method.

### Description

#### Reference





10





### APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com