

Recombinant Bovine Monokine Induced by Interferon-gamma/CXCL9

Information

Gene ID	513990
Accession #	A9QWP9
Alternate Names	
Source	Escherichia coli.
M.Wt	Approximately 11.9 kDa, a single non-glycosylated polypeptide chain containing 104 amino acids. But it migrates with an apparent molecular mass of 18 kDa in SDS-PAGE.
AA Sequence	VPAIRNGRCS CINTSQGMIH PKSLKDLKQF APSPSCEKTE IIATMKNGNE ACLNPDLP EV KELIKEWEKQ VNQKKKQKRG KKYKKTKKVP KVKRSQRPSQ KKT
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.0, 500 mM NaCl.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20°C. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human lymphocytes is in a concentration range of 0.1-1.0 ng/ml.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	10µg	100µg	500µg
Recombinant Bovine Monokine Induced by Interferon-gamma/CXCL9	10µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

Quality Control

Purity	> 96 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 0.1 EU/ μ g of rBoMIG/CXCL9 as determined by LAL method.

Description

CXCL9 is a T-cell chemoattractant induced by IFN- γ belonging to the CXC chemokine family and it is also known as Monokine induced by gamma interferon (MIG). CXCL9 is closely related to two other CXC chemokines called CXCL10 and CXCL11 and they all elicit their chemotactic functions by interacting with the chemokine receptor CXCR3. CXCL9 is a cytokine that affects the growth, movement, or activation state of cells that participate in immune and inflammatory response and chemotactic for activated T-cells. Recombinant bovine CXCL9 contains 104 amino acids which is a single non-glycosylated polypeptide chain. The bovine CXCL9 shares 92 % and 72 % a.a. sequence identity with sheep and human CXCL9.

Reference

APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com