

Recombinant Murine Desert Hedgehog Homolog Cys23llelle

Information

oximately 20.0 kDa, a single non-glycosylated polypeptide chain containin amino acids. GRGPVG RRRYVRKQLV PLLYKQFVPS MPERTLGASG PAEGRVTRGS RDLVPNY NPDIIFKDEE NSGADRLMTE RCKERVNALA IAVMNMWPGV /TEGWDE DGHHAQDSLH YEGRALDITT SDRDRNKYGL LARLAVEAGF YYESRNH IHVSVKADNS LAVRAGG e Filtered White lyophilized (freeze-dried) powder. a manual defrost freezer and avoid repeated freeze-thaw cycles nonths from date of receipt, -20 to -70 °C as supplied onth, 2 to 8 °C under sterile conditions after reconstitution onths, -20 to -70 °C under sterile conditions after reconstitution	
oximately 20.0 kDa, a single non-glycosylated polypeptide chain containin amino acids. GRGPVG RRRYVRKQLV PLLYKQFVPS MPERTLGASG PAEGRVTRGS RDLVPNY NPDIIFKDEE NSGADRLMTE RCKERVNALA IAVMNMWPGV /TEGWDE DGHHAQDSLH YEGRALDITT SDRDRNKYGL LARLAVEAGI YYESRNH IHVSVKADNS LAVRAGG e Filtered White lyophilized (freeze-dried) powder. a manual defrost freezer and avoid repeated freeze-thaw cycles nonths from date of receipt, -20 to -70 °C as supplied onth, 2 to 8 °C under sterile conditions after reconstitution onths, -20 to -70 °C under sterile conditions after reconstitution	
amino acids. GRGPVG RRRYVRKQLV PLLYKQFVPS MPERTLGASG PAEGRVTRGS RDLVPNY NPDIIFKDEE NSGADRLMTE RCKERVNALA IAVMNMWPGV /TEGWDE DGHHAQDSLH YEGRALDITT SDRDRNKYGL LARLAVEAGE YYESRNH IHVSVKADNS LAVRAGG e Filtered White lyophilized (freeze-dried) powder. a manual defrost freezer and avoid repeated freeze-thaw cycles months from date of receipt, -20 to -70 °C as supplied onth, 2 to 8 °C under sterile conditions after reconstitution onths, -20 to -70 °C under sterile conditions after reconstitution hilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4, 1 mM	
RDLVPNY NPDIIFKDEE NSGADRLMTE RCKERVNALA IAVMNMWPGV /TEGWDE DGHHAQDSLH YEGRALDITT SDRDRNKYGL LARLAVEAG YYESRNH IHVSVKADNS LAVRAGG e Filtered White lyophilized (freeze-dried) powder. a manual defrost freezer and avoid repeated freeze-thaw cycles nonths from date of receipt, -20 to -70 °C as supplied onth, 2 to 8 °C under sterile conditions after reconstitution onths, -20 to -70 °C under sterile conditions after reconstitution nilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4, 1 mM	
a manual defrost freezer and avoid repeated freeze-thaw cycles nonths from date of receipt, -20 to -70 °C as supplied onth, 2 to 8 °C under sterile conditions after reconstitution onths, -20 to -70 °C under sterile conditions after reconstitution nilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4, 1 mM	
nonths from date of receipt, -20 to -70 °C as supplied onth, 2 to 8 °C under sterile conditions after reconstitution onths, -20 to -70 °C under sterile conditions after reconstitution nilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4, 1 mM	
Lyophilized from a 0.2 μ m filtered concentrated solution in PBS, pH 7.4, 1 mM DTT, 0.05 % Tween-80.	
ecommend that this vial be briefly centrifuged prior to opening to bring the ents to the bottom. Reconstitute in sterile distilled water or aqueous buffer ining 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions Id be apportioned into working aliquots and stored at \leq -20 °C. Further ons should be made in appropriate buffered solutions.	
Fully biologically active when compared to standard. The ED as determined by its ability to induce alkaline phosphatase production by murine MC3T3-E1 cells is 5-20 μ g/ml.	
Gel pack.	
Centrifuge the vial prior to opening.	
Research Use Only! Not to be used in humans.	

Components and Storage

Components	5µg	100µg	500µg
Recombinant Murine Desert Hedgehog Homolog Cys23llelle	5µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

l	Quality Control		B10	
	Purity	> 98 % by SDS-PAGE and HPLC analyses.	Recommended	
	Endotoxin	Less than 0.1 EU/ μ g of rMuDHH C23II as determined by LAL method.		

Description

Reference



APER BIO





