

Anti-CD63 Rabbit Polyclonal Antibody

Introduction

Functions as cell surface receptor for TIMP1 and plays a role in the activation of cellular signaling cascades. Plays a role in the activation of ITGB1 and integrin signaling, leading to the activation of AKT, FAK/PTK2 and MAP kinases. Promotes cell survival, reorganization of the actin cytoskeleton, cell adhesion, spreading and migration, via its role in the activation of AKT and FAK/PTK2.

Product parameters

Alternative Names	CD63; MLA1; TSPAN30; CD63 antigen; Granulophysin; Lysosomal-associated membrane protein 3 LAMP-3; Melanoma-associated antigen ME491; OMA81H; Ocular melanoma-associated antiger Tetraspanin-30; Tspan-30; CD antigen CD63
Gene ID	967
Gene Name	CD63
SwissProt ID	P08962
Host	Rabbit A PEX B O
Reactivity	Human
Molecular Weight	Calculated MW: 26 kDa; Observed MW: 26 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Polyclonal Antibody
Clonality No.	-
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human CD63.
Purification	Affinity Purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	WB, IHC-F, IHC-P, ICC/IF, ELISA
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 ELISA: 1/10000
Research Field	Cardiovascular

Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.





Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





