

Anti-MyD88 Rabbit Polyclonal Antibody

Introduction

Members of the Toll-like receptor (TLR) family, named for the closely related Toll receptor in *Drosophila*, play a pivotal role in innate immune responses. TLRs recognize conserved motifs found in various pathogens and mediate defense responses. Triggering of the TLR pathway leads to the activation of NF- κ B and subsequent regulation of immune and inflammatory genes.

Product parameters

Alternative Names	MYD88; Myeloid differentiation primary response protein MyD88
Gene ID	4615
Gene Name	MYD88
SwissProt ID	Q99836
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 33 kDa; Observed MW: 33 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Polyclonal Antibody
Clonality No.	-
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	The antiserum was produced against synthesized peptide derived from human MyD88.
Purification	Affinity Purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	WB, IHC-F, IHC-P, ICC/IF, ELISA
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 ELISA: 1/10000
Research Field	Signal Transduction
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol



Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

