

Anti-Transcriptional Enhancer Factor TEF 1 Rabbit Polyclonal Antibody Introduction

Transcription factor which plays a key role in the Hippo signaling pathway, a pathway involved in organ size control and tumor suppression by restricting proliferation and promoting apoptosis.

Product parameters

| Alternative Names | TEAD1; TCF13; TEF1; Transcriptional enhancer factor TEF-1; NTEF-1; Protein GT-IIC; TEA domain family member 1; TEAD-1; Transcription factor 13; TCF-13 |
|--------------------|--|
| Gene ID | 7003 |
| Gene Name | TEAD1 |
| SwissProt ID | P28347 |
| Host | Rabbit |
| Reactivity | Human, Mouse, Rat |
| Molecular Weight | Calculated MW: 48 kDa; Observed MW: 52 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Polyclonal Antibody |
| Clonality No. | - |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Free |
| Immunogen | A synthesized peptide derived from human TEAD1 |
| Purification | Affinity Purified |
| Buffer System | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Application | WB, IHC-P, IP |
| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/50 |
| Research Field | Epigenetics and Nuclear Signaling |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |

| Expiration Date | 12 months |
|-----------------|----------------------------------|
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





