

## Mouse IgG Isotype Control

# Introduction



Isotype control antibodies are used to estimate the nonspecific binding of target primary antibodies due to Fc receptor binding or other protein-protein interactions. An isotype control antibody should have the same immunoglobulin type and be used at the same concentration as the Target antibody.

### Product parameters

Alternative Names	-
Gene ID	-
Gene Name	Mouse IgG
SwissProt ID	-
Host	Mouse
Reactivity	Isotype Control
Mole <mark>cular</mark> Weight	*BIO APExBIO
Conjugation	Unconjugated Administration Explore the Velocom
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Polyclonal Antibody
Clonality No.	-
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	-
Purification	Affinity Purified
Buff <mark>er S</mark> ystem	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	FC, IP, ChIP
Dilution Ratio	IP: 1/20 FC: 1/100 ChIP: 1/20
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Shipping	Blue ice

Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

### **Protocol**

Configure the product according to the application range and recommended dilution ratio.



\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

#### Note

This product is for scientific research use only.





