

Recombinant Human EGF

Accession #	P01133
Alternate Names	beta-urogastrone; EGF; epidermal growth factor (beta-urogastrone); epidermal growth factor; hEGF
Source	Human embryonic kidney cell, HEK293-derived human EGF protein
Protein sequence	Asn971-Arg1023
M.Wt	6.3 kDa
Appearance	Solution protein.
Stability & Storage	Avoid repeated freeze-thaw cycles. It is recommended that the protein be aliquoted for optimal storage. 3 years from date of receipt, -20 to -70 °C as supplied.
Concentration	0.2 mg/mL
Formulation	Dissolved in sterile PBS buffer.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. This solution can be diluted into other aqueous buffers.
Biological Activity	Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED50 for this effect is 4-50 pg/mL.
Shipping Condition	Shipping with dry ice.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Quality Control

Purity	> 95%, determined by SDS-PAGE.
Endotoxin	<0.010 EU per 1 ug of the protein by the LAL method.

Description

Epidermal growth factor (EGF) is a small, potent growth factor capable of inducing cell proliferation, differentiation, and survival. EGF is the founding member of the EGF family that also includes TGF- α , amphiregulin (AR), betacellulin (BTC), epiregulin (EPR), heparin-binding EGF-like growth factor (HB-EGF), epigen, and the neuregulins (NRG)-1 through -6 [1]. Members of The EGF family are characterized by a shared structural motif, the EGF-like domain, which contains three intramolecular disulfide bonds that are formed by six similarly spaced, conserved cysteine residues [2]. These disulfide bonds are essential for proper protein conformation and receptor binding. All EGF family members are synthesized as type I transmembrane precursor proteins that may contain several EGF domains in the extracellular region. The mature proteins are released from the cell surface by regulated proteolysis [1]. The full length EGF protein is 1207 amino acids (aa) (EGF precursor)

containing nine EGF domains and nine LDLR class B repeats. However, the mature protein is much smaller, only 53 aa, and is generated by proteolytic cleavage of the EGF domain proximal to the transmembrane region [3]. EGF is well conserved across mammals with mature human EGF 70% identical to mature mouse and rat EGF. Physiologically, EGF is found in various body fluids, including blood, milk, urine, saliva, seminal fluid, pancreatic juice, cerebrospinal fluid, and amniotic fluid [4]. EGF is a high affinity ligand of the EGF receptor (ErbB). Four ErbB (HER) family receptor tyrosine kinases including EGFR/ErbB1, ErbB2, ErbB3 and ErbB4, mediate responses to EGF family members [5]. EGF binding induces dimerization of the EGF receptor resulting in activation of the protein tyrosine kinase signaling pathway. These receptors undergo a complex pattern of ligand-induced homo- or hetero-dimerization to transduce EGF family signals [6, 7].

Reference

- [1]. Harris, R.C. et al. (2003) Exp. Cell Res. 284:2.
- [2]. Carpenter, G. and Cohen, S. (1990) J. Biol. Chem. 265:7709.
- [3]. Bell, G.I. et al. (1986) Nucl. Acids Res. 14:8427.
- [4]. Carpenter, G. and Zengdegui, J.G. (1986) Exp. Cell Res. 164:1.
- [5]. Jorissen, R.N. et al. (2003) Exp. Cell Res. 284:31.
- [6]. Gamett, D.C. et al. (1997) J. Biol. Chem. 272:12052.
- [7]. Qian, X. et al. (1994) Proc. Natl. Acad. Sci. 91:1500.

APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com